

The heat flux from a relativistic kinetic equation with a simplified collision kernel

A. Sandoval-Villalbazo*, A. L. Garcia-Perciante[†] and L.S. Garcia-Colin**

**Depto. de Fisica y Matematicas, Universidad Iberoamericana, Prolongacion Paseo de la Reforma 880, Mexico D. F. 01219, Mexico.*

[†]*Depto. de Matematicas Aplicadas y Sistemas, Universidad Autonoma Metropolitana-Cuajimalpa, Artificios 40 Mexico D.F 01120, Mexico.*

***Depto. de Fisica, Universidad Autonoma Metropolitana-Iztapalapa, Av. Purisima y Michoacan S/N, Mexico D. F. 09340, Mexico. Also at El Colegio Nacional, Luis Gonzalez Obregon 23, Centro Historico, Mexico D. F. 06020, Mexico*

Abstract. We show how using a special relativistic kinetic equation with a BGK- like collision operator the ensuing expression for the heat flux can be casted in the form required by Classical Irreversible Thermodynamics. Indeed, it is linearly related to the temperature and number density gradients and not to the acceleration as the so-called "first order in the gradients theories" contend. Here we calculate explicitly the ensuing transport coefficients and compare them with the results obtained by other authors.

Keywords: Relativistic fluid dynamics, Relativistic kinetic theory, Heat transfer

PACS: 47.75.+f, 05.70.Ln, 51.30.+y

The purpose of this work is to calculate the explicit form for the heat flux in an ideal relativistic gas to find out what is the form of the constitutive equation as required by Classical Irreversible Thermodynamics (CIT), in the temperature and particle number density representation. This result is important by itself because it will help in clarifying whether or not such constitutive equation should contain a term proportional to the acceleration as it has been claimed by many authors. To accomplish this task, we start by considering the simplified relativistic Boltzmann equation in the absence of external forces:

$$v^\alpha f_{,\alpha} = J(f, f') \quad (1)$$

The term on the right hand side of Eq. (1) is the collision operator which is modeled by means of the BGK approximation [1]. In Eq. (1), the molecular four velocity, v^α is

$$v^\alpha = \begin{bmatrix} \gamma w^\ell \\ \gamma c \end{bmatrix} \quad (2)$$

where w^ℓ is the molecular three-velocity and $\gamma = (1 - w^\ell w_\ell / c^2)^{-1/2}$ is the usual relativistic factor. All greek indices run from 1 to 4 and the latin ones run up to 3. The number density reads

$$n = \int f \gamma dv^* \quad (3)$$

with $dv^* = \gamma^5 \frac{cd^3w}{v^4}$ [2]. Now, standard kinetic theory leads to the energy balance equation [1, 3]:

$$\frac{\partial}{\partial t} (n \langle mc^2 \gamma \rangle) + \left(n \langle mc^2 \gamma w^\ell \rangle \right)_{;\ell} = 0 \quad (4)$$

where the thermodynamic average is defined through:

$$\langle \psi \rangle = \frac{1}{n} \int \gamma \psi f dv^* \quad (5)$$

In Eq. (4) one can easily identify the heat flux as:

$$J_{[Q]}^\ell = n \langle mc^2 v^\ell \rangle = mc^2 \int v^\ell f \gamma dv^* \quad (6)$$

In a Chapman-Enskog expansion, the distribution function is written as

$$f = f^{(0)} (1 + \phi) \quad (7)$$

where ϕ is the first order correction in the Knudsen parameter, a weighted measure of the gradients in the system. For particles of rest mass m , relativistic parameter $z = \frac{kT}{mc^2}$ and in the non-degenerate case, the equilibrium function reads [1, 3]:

$$f^{(0)} = \frac{n}{4\pi c^3 z \mathcal{K}_2\left(\frac{1}{z}\right)} e^{\frac{u^\beta v_\beta}{zc^2}} \quad (8)$$

in which k is Boltzmann's constant, u^β the hydrodynamic four-velocity and $\mathcal{K}_2$ is the modified Bessel function of the second kind. We remind the reader that the first order correction ϕ contains already the dissipative effects in the system.

If τ denotes the collisional time appearing in the BGK model, using the functional hypothesis [4], the function ϕ in Eq. (7) is given in terms of the thermodynamic forces as

$$\phi = -\tau v^\alpha \left(\frac{\partial f^{(0)}}{\partial n} n_{,\alpha} + \frac{\partial f^{(0)}}{\partial T} T_{,\alpha} + \frac{\partial f^{(0)}}{\partial u^\beta} u_{,\alpha}^\beta \right) \quad (9)$$

where the derivatives of the Jüttner distribution function are

$$\frac{\partial f^{(0)}}{\partial n} = \frac{f^{(0)}}{n} \quad (10)$$

$$\frac{\partial f^{(0)}}{\partial T} = \left(-1 - \frac{\mathcal{K}_1\left(\frac{1}{z}\right)}{2z\mathcal{K}_2\left(\frac{1}{z}\right)} - \frac{\mathcal{K}_3\left(\frac{1}{z}\right)}{2z\mathcal{K}_2\left(\frac{1}{z}\right)} \right) \frac{f^{(0)}}{T}, \quad (11)$$

and

$$\frac{\partial f^{(0)}}{\partial u^\beta} = \frac{v_\beta}{zc^2} f^{(0)} \quad (12)$$




FIGURE 1. The functions $f_T(z)$ (left) and $f_n(z)$ (right).

We now introduce Euler's equations for the time derivatives in (9). A hydrodynamic acceleration term appears in the last term of Eq. (9) for $\alpha = 4$. In the standard approach this term is expressed by the pressure gradient using Euler's equation. This would lead to an awkward representation involving n , T , p and u^α . However, in order to comply with the requirements of CIT, using the fact that the ordinary non-equilibrium state variables of the fluid are n , u^α and T , we substitute the pressure gradient through the equation of state $p = nkT$, which readily leads to the following form for the heat flux

$$J_{[Q]}^\ell = -L_{TT} \frac{T^{\ell}}{T} + L_{nT} \frac{n^{\ell}}{n} \quad (13)$$

Equation (13) is of the canonical form required by CIT, the thermodynamic forces are the temperature and the density gradients, respectively. The transport coefficients are given by

$$L_T = \frac{nk^2T^2}{m} \tau f_T(z), \quad L_n = \frac{nk^2T^2}{m} \tau f_n(z) \quad (14)$$

where the functions f_T and f_n are given by

$$f_T(z) = \left(\frac{1}{z} - \left(4z + \frac{\mathcal{K}_1(\frac{1}{z})}{\mathcal{K}_2(\frac{1}{z})} \right)^{-1} \right) \left(\frac{1}{z} + 5 \frac{\mathcal{K}_3(\frac{1}{z})}{\mathcal{K}_2(\frac{1}{z})} \right) - \left(1 + \frac{\mathcal{K}_1(\frac{1}{z})}{2z\mathcal{K}_2(\frac{1}{z})} + \frac{\mathcal{K}_3(\frac{1}{z})}{2z\mathcal{K}_2(\frac{1}{z})} \right) \frac{\mathcal{K}_3(\frac{1}{z})}{z\mathcal{K}_2(\frac{1}{z})} \quad (15)$$

$$f_n(z) = \left(4z + \frac{\mathcal{K}_1(\frac{1}{z})}{\mathcal{K}_2(\frac{1}{z})} \right)^{-1} \left(\frac{1}{z} + 5 \frac{\mathcal{K}_3(\frac{1}{z})}{\mathcal{K}_2(\frac{1}{z})} \right) - \frac{\mathcal{K}_3(\frac{1}{z})}{z\mathcal{K}_2(\frac{1}{z})} \quad (16)$$

and are shown in Figure 1.

The coefficient given in Eq. (16) has not been analyzed before, although it has been proposed in Ref. [5]. Its presence in the constitutive equation for the heat flux given in Eq. (13), which has the structure required by CIT, guarantees the validity of Onsager's

regression of fluctuations hypothesis. Other authors obtain an expression for the heat flux in terms of the acceleration or pressure gradient which is, not only against the tenets of CIT, but also gives rise to the so-called generic instabilities for the relativistic fluid [6, 7]. In contrast, our approach provides a picture in which fluctuations do not grow exponentially within the first order in the gradients description of the relativistic simple fluid.

Two important facts shown in Figure 1 are worth pointing out as a final remark. Firstly, we have obtained an analytical expression for the thermal conductivity for the relativistic gas, using in the BGK approximation. This result is new and differs from previous calculations since this is the first time the representation given in Eq. (13) is explicitly used. Notice also how this coefficient yields the correct non-relativistic limit $L_T \sim \frac{5}{2} \frac{nk^2 T^2}{m} \tau$. Secondly, the figure on the right shows a novel result, namely a coefficient relating a heat flux to a number density gradient present only in the relativistic case. This coefficient vanishes for $z \rightarrow 0$.

REFERENCES

1. C. Cercignani y G. Medeiros Kremer; *The relativistic Boltzmann equation: theory and applications*, Birkhäuser, Berlin (2002).
2. R. L. Liboff y R. C. Liboff; *Kinetic theory: classical, quantum and relativistic descriptions*, Springer, Berlin (2003).
3. S. R. de Groot, W. A. van Leeuwen & Ch. G. van Weert; *Relativistic Kinetic Theory*, North Holland, Netherlands (1980).
4. G. E. Uhlenbeck & G. W. Ford, *Lectures in Statistical Mechanics*, American Mathematical Society, Providence RI (1963).
5. A. Sandoval-Villalazo, A. L. Garcia-Perciante and L. S. Garcia-Colin, arXiv: 0805.4237 (2008).
6. A. L. Garcia-Perciante, L. S. Garcia-Colin and A. Sandoval-Villalazo, *Gen. Rel. and Grav.* (2009, in press) arXiv:0802.3838.
7. W. A. Hiscock and L. Lindblom, *Phys. Rev. D* **31**, 725 (1985).